Global J. Environ. Sci. Manage. 5(2): 225-236, Spring 2019

Global Journal of Environmental Science and Management
(GJESM)

Homepage: https://www.gjesm.net/

ORIGINAL RESEARCH PAPER

Contamination profile of heavy metals in marine fish and shellfish

M.A. Salam[1,2,*]; S.C. Paul[3]; S.N.B.M. Noor[2]; S.A. Siddiqua[4]; T.D. Aka[5]; R. Wahab[6]; E.R. Aweng[2]

[1]Department of Environmental Science and Disaster Management, Noakhali Science and Technology University, Noakhali, Sonapur -3814, Bangladesh

[2]Faculty of Earth Science, University, University Malaysia Kelantan, Jeli Campus, 17600 Jeli, Malaysia

[3]Department of Applied Chemistry and Chemical Engineering, Noakhali Science and Technology University, Noakhali, Sonapur -3814, Bangladesh

[4]Department of Animal Breeding and Genetics, Bangladesh Agricultural University, Mymensingh- 2202, Bangladesh

[5]Department of Pharmacy, Noakhali Science and Technology University, Noakhali, Sonapur -3814, Bangladesh

[6]School of Engineering and Technology, University College of Technology Sarawak, Malaysia.

ARTICLE INFO

Article History:
Received 21 October 2018
Revised 13 February 2019
Accepted 21 February 2019

Keywords:
Crab
Estimated Daily Intake (EDI)
Fish
Perak River
Prawn
Target Hazard Quotient (THQ)

ABSTRACT

Rapid industrialization along with advanced agricultural activities led to the contamination in aquatic environment with heavy metals. Heavy metals ultimately pass into human body through having aquatic animals like fish, prawn and crab. In this study, accumulation of heavy metals (zinc, copper, iron, cadmium and lead) in various organs of four commonly consumed fish (*Euthynnus affinis*, *Pampus argenteus, Descapterus macrosoma,* and *Leiognathus daura*), prawn (*Fenneropenaeus indicus) and* crab *(Portunus pelagicus)* of Tok Bali Port, Kelantan, Malaysia were determined. Health risk was assessed using estimated daily intake and target hazard quotients. Although the concentrations of all the heavy metals in all fish, prawn and crab species were lower as per Malaysian Food Act, but the concentrations showed remarkable differences among the species and organs. The concentration of heavy metals in the gill was the highest of all fish species followed by in the liver and flesh. The total accumulation of heavy metals was maximum in *Euthynnus affinis* followed by *Leiognathus daura*, *Descapterus macrosoma* and *Pampus argenteus* of the fish species. However, the highest concentrations (µg/g) was for Zn (72.97±2.75), followed by Fe (4.309±0.68), Cd (1.189±0.78), Cu (1±0.87) and Pb (0.41±0.19) among all the heavy metal contents of fish. No significant variation ($P<0.05$) of the heavy metal concentration in prawn and crab species was observed. The investigation indicated that the fish, prawn and crab species of this port were safe for human consumption but the safe disposal of various wastes should be practiced to control the heavy metal accumulation in future.

DOI: 10.22034/gjesm.2019.02.08

INTRODUCTION

Fish is consumed all over the world for its nutritional value, along with its high quality proteins, great omega-3 fatty acid content, and low saturated fat content and due to having a good level of vitamin-B (Özden *et al.,* 2018; Hajeb *et al.,* 2009; Arulkumar *et al.*, 2017; Gu *et al.,* 2017). It is anticipated that fish contributes about 17% of animal protein and almost 6% of all protein consumed by human beings (Jos´e *et al.,* 2007). However fish become threat to human health because of heavy metals accumulation in marine and fresh water via various urbanization,

*Corresponding Author:
Email: *s_salam1978@yahoo.com*
Phone: +88 0312 61433
Fax: 321 62788

Note: Discussion period for this manuscript open until July 1, 2019 on GJESM website at the "Show Article.

agricultural and industrial activities (Li *et al.,* 2017; Martin *et al.,* 2015; Zhang *et al.,* 2011). Heavy metals like Cadmium (Cd), Chromium (Cr), Lead (Pb), Iron (Fe), Manganese (Mn), Copper (Cu), Nickel (Ni), Arsenic (As), Mercury (Hg) and Zinc (Zn) have high level of toxicity, and persistence capacity possessing potential for biomagnification, bioaccumulation and incorporation into the food chain after reaching a certain limit in the aquatic environment (Hosseini *et al.,* 2012; Zhang *et al.,* 2010; Ahmed *et al.,* 2015; Kibria *et al.,* 2016). Bioaccumulation and biomagnification denote the processes and pathways of heavy metal pollutants from one trophic level to others in the food-web (Part *et al.,* 1985). In the aquatic environment heavy metals are assimilated by fish, since fish is great bioaccumulator, assimilation can occur by ingestion, ion exchange across gills or membrane surface and adsorption by tissues of the fish (Ahmed *et al.,* 2014). Fish also have diverse bioaccumulation characteristics, and accumulation can occur in the head, liver, bones, kidney, stomach, heart, muscle, operculum, vertebrae and flesh of fish species *(Murtala et al., 2012; Zhao et al.,* 2012). It happens in different parts of different fish species due to the varying soluble nature of metals in water, bioavailability and the different habitats, life cycles, nature of feeding, ecology and physiological nature of fish (Perugini *et al.,* 2014; Anandkumar *et al.,* 2017). High bioaccumulation of heavy metals in the aquatic body can lead to genotoxic damage of aquatic species and high concentrations of heavy metals can have cytotoxic, mutagenic and genotoxic effects on fish species (Matos *et al.,* 2017). Consumption of fish with high level of heavy metal contamination can be a serious health hazard (Salamat *et al.,* 2015). Cd can be seriously injurious for health, causing kidney damage, respiratory dysfunction, renal and hepatic system damage, bone disease, hyper tension, poor reproductive capacity, tumors etc. due to its long term uptake by the human body (Satarug *et al.,* 2017; Lee *et al.,* 2011). The accumulation of Hg in the human body causes damage to distinct visual cortex areas, and neuronal loss of adult brain (Al-Busaidi *et al.,* 2011) and neural impairment in infants (Harada *et al.,* 1994). Pb can have a contribution to renal failure as well as liver damage in humans (Lee *et al.,* 2011). Long-term Pb consumption can cause mental retardation and even mortality of humans (Wani *et al.,* 2015). Non- essential heavy metals (Cd and Pb) even at trace amounts can exhibit extreme toxicity while essential metals (Cu, Zn and Fe) can also cause health problems due to overdosing. As heavy metals cannot be degraded they accumulate in aquatic organisms or animal and are deposited in sediments (Merian 1991, Linnik *et al.,* 2000; Gale *et al.,* 2004, Wei *et al.,* 2014), it is very essential to assess the health risk assessment of heavy metals in human health through consumption of aquatic animal. The sources of heavy metals in water-bodies include the paint on ships, diverse industrial processes e.g. stabilizers for PVC and agricultural activities such as the use of fungicides, pesticides etc. (Lee *et al.,* 2016). Where rapid industrialization and mechanized agricultural activities enhance the heavy metal pollution rate the environmental hazards to invertebrates, and vertebrates,- including fish and humans are increased (Bashir *et al.,* 2013; Yi *et al.,* 2017). Fast moving industrialization and economic development in Malaysia has led to the utilization of toxic chemicals including heavy metals, and has resulted in increasing water pollution in its coastal regions (Bashir *et al.,* 2013, Agusa *et al.,* 2005). Tok Bali Port is the largest port in Kelantan and makes a great economic contribution to Malaysia. However, the various shipping and transportation activities around this port may also contribute to heavy metal pollution in sea water and subsequent accumulation in commercially important marine fishes, prawn and crabs (Salleh and Halim 2018). Several studies in port city and coastal region around the world shows the presence of heavy metals in commercially important fish, prawn and crab species and their health effects (Amisah *et al.,* 2011; Schintu *et al.,* 2016; Sany *et al.,* 2012; Nemr *et al.,* 2016; Sharif *et al.,* 1993; Velusamy *et al.,* 2014; Gu *et al.,* 2015; Sarkara *et al.,* 2016). However there is less data available about the presence of heavy metals in fish, prawn and crabs consumed around the Malaysia and consequently about their contribution to the dietary intake of the Malaysian population. Considering this situation it is important to determine the concentration of heavy metals in edible commercially important fish, prawn and crab species available in the east coast region of Tok Bali Port, Kelantan, Malaysia. In this study concentration of heavy metals (Zn, Cu, Fe, Cd and Pb) in various organs of four commonly consumed fish namely *Euthynnus affinis (*eastern little tuna), *Pampus argenteus* (silver Pomfret)*, Decapterus macrosoma* (Layang scad), and *Leiognathus daura* (blackfin ponyfish), one prawn species namely *Fenneropenaeus indicus* (Indian white prawn) and one crab species *Portunus pelagicus* (blue crab) were determined along with various health risk

assessment indexes (EDI and THQ). The study has been carried out in Malaysia in 2015.

MATERIALS AND METHODS

Study area

Tok Bali Port is the largest port in Kelantan, Malaysia and provides a major food source to the community of Kelantan. Tok Bali is located about 48 kilometer from Kota Bharu and is considered as a major player in marine services in the state of Kelantan. The shipping and transportation industry near to this port are major sources of water pollution. Greywater from ships also contributes to the remarkable pollution in this port. Fish, prawn and crab samples were collected from Tok Bali Port in Pasir Puteh, Kelantan with an exact location of $5^0$53'49.1"North and $102^0$29'06.8"Eeast in 2015. The sampling location is given in Fig. 1.

Collection, sample preparation and heavy metals analysis

Four fish species, namely, layang scad (n=10), estern little tuna (n=10), blackfin ponyfish (n=10), and silver pomfret (n=10) and two different species of shellfishes namely, indian white prawn (n=10), and blue crab (n=10) were collected in January, 2016. The size and weights of the collected fish and shellfish are given in Table 1. The samples were primarily retained in the zip-locked acid-washed polyethylene bags at -20 °C and finally transported to the laboratory for further processing. In the laboratory, after separating the gills, livers and muscles the sample tissue was homogenized by grinding. The homogenized sample was then freeze dried at (-10 °C) and ground to a fine powder. After adding 5 ml of concentrated nitric acid and 5 ml of hydrogen peroxide to the 5 g of dried sample in the digester and stored at room temperature for 24 h (Asegbeloyin *et al.,* 2010). Finally, the concentration of heavy metals were determined by atomic absorption spectrometer (AAS); Perkin-Elmer-3300 by using standard diluted sample solution. For each set of experiments blanks, certified reference materials (CRMs), and samples were run and corrections applied where necessary. All of the experiments were carried out in triplicate to eliminate any batch-specific errors and only average values were reported (Islam *et al .,* 2014). All laboratory equipment used was washed with




Fig. 1. Geographic location of the study area along with the sampling sites in Kelantan, Malaysia

Table 1. List of fish, prawn and crab species with length, weight and Fulton's condition factor (K) along with ecological characteristics

Fish species					
English name	Scientific name	Habitat	Length (cm)	Weight (g)	K
Layang Scad	*Descapterus macrosoma*	Pelagic (reef slopes)	15.5	57.5	1.54
Estern little tuna	*Euthynnus affinis*	Epi-pelagic	25.1	181.01	1.14
Blackfin ponyfish	*Leiognathus daura*	Demersal and pelagic (water column)	32.01	509.98	1.56
Silver pomfret	*Pampus argenteus*	Benthopelagic	23.5	164.05	1.26
Prawn					
Indian white prawn	*Fenneropenaeus indicus*	Demersal	-	200	-
Crab					
Blue crab	*Portunus pelagicus*	Intertidal	-	200	-

phosphate-free soap, double rinsed with distilled water and left in 10% HNO_3 for 24 h to prevent contaminations. Calibration, the use of blanks, spike samples, the performance characteristics of the procedure and the reporting of outcome were the significant aspect of QA and QC (Natural Resources Management and Environment Department, 1998) in the study. Recovery of all metals studied ranged from 90 to 98.6%.

Statistical analysis

Descriptive statistical analysis throughout the study was carried out by using Statistical Package for the Social Sciences (SPSS) 16 software. The statistical significance of the heavy metals concentrations within the same tissue and the heavy metals concentrations among the six different species were calculated by using the Tukey (Honestly Significant Difference) HSD test. Correlation values among heavy metals content among various organs and fish length were calculated by SPSS. Fulton condition factor (K) was calculated (Table 1) according to Eq. 1 (Htun-Han, 1978).

$$K = 100\left(\frac{W}{L^3}\right) \quad (1)$$

Where W is the weight of fish in gram (gm) and L is the length of fish in centimeter (cm).

Health risk assessments

The estimated daily intake of metals was estimated by using Eq. 2 (Onsanit *et al.,* 2010).

Estimated daily intake (EDI)=

$$\frac{metal\ concetation\left(\frac{ug}{g}w\ /w\right)\times consumtion\ rate\left(\frac{g}{d}\right)}{body\ weight\ (kg)} \quad (2)$$

Metals concentration in the fish, prawn and crab used in above equation, were the average maximum values of trace element concentrations in muscle. The body weights for children and adults were assumed to be 32.7 kilogram (kg) and 55.9 kg, respectively, the consumption rate was 50 g/day and 93 g/day for children and adults (Zhao *et al.,* 2012). The Tolerable intake was considered in this study to examine the EDI of metals in fishes, prawns and crabs. The tolerable intake for Zn, Cu, Fe Cd and Pb are 300-1000, 50-500, 80-100, 1 and 3.57 µg/kg/day, respectively as Eq. 3. (JECFA, 2009).

$$\text{Target hazardous quotient (THQ)} = \frac{\text{EF X ED X FIR X C}}{RfD\ X\ WAB\ X\ TA}10^{-3} \quad (3)$$

Here EF represents exposure frequency in day/year; ED represents exposure duration (70 years and 6 years for adult and children respectively); FIR (Food ingestion rate) represents the food ingestion rate (93 g/Day and 50 g/Day for adults and children); C is the measured concentration; RfD = reference dose (µg/g); WAB is average body weight in kg and TA is average exposure time for non-carcinogens (365 day/year X ED). There would be no obvious risk if the calculated THQ is less than 1 (USEPA, 2011).

RESULTS AND DISCUSSION

Physical characteristics of fish, prawn and crab

Table 1 shows the values of Fulton's condition factor calculated in this study for all analyzed samples. The value of "k" greater than one suggesting that well grown fish were collected and also indicating that the fish were in good condition (Nasri *et al.,* 2017).

Comparisons of heavy metals in selected fish, prawn and crab species

Age, diet, metabolism, size and length of the fish

should be considered as the factors for analyzing the fish species of same coastal environment (Noelia *et al.,* 2017). The concentrations of five heavy metals (Zn, Cu, Fe, Cd and Pb) have been determined in the tissues (i.e. gill, liver and flesh) of four fish species, one prawn and one crab species in the coastal area of Tok Bali, Kelatan, Malaysia are listed in Table 2. The ranges of concentration of heavy metal content are as follows: 72.970 - 4.827 µg/g for Zn, 0.789 - 0.018 µg/g for Cu, 4.309 - 0.033 µg/g for Fe, 1.189 -0.003µg/g for Cd and 0.410 - 0.001 µg/g for Pb. The highest concentrations was for Zn, followed by Fe, Cd, Cu and Pb. The order of species according to the heavy metal accumulation is as follows: *Leiognathus daura> Euthynnus affinis> Descapterus macrosoma> Pampus argenteus* for Zn, *Euthynnus affinis> Pampus argenteus> Descapterus macrosoma>Leiognathus daura* for Cu, *Leiognathus daura> Euthynnus affinis> Descapterus macrosoma> Pampus argenteus* for Fe, *Euthynnus affinis> Leiognathus daura> Pampus argenteus> Descapterus macrosoma* for Cd and Pb, as seen in Table 2. In the case of prawn and crab species the highest amounts of Zn and Fe were found in *Fernneropenaeus indicus* 14.40 µg/g and 2.11µg/g respectively because *Fernneropenaeus indicus* (indian white prawn) are usually prefers mud or sandy mud at depths of 2 - 90 meters. The highest concentrations of Cu, Cd and Pb were 1.0 µg/g, 0.1µg/g and 0.02µg/g respectively and were found in *Portunus pelagicus*. Among all species of fish, prawn and crab, the hazardous heavy metal content (Pb and Cd) in shellfish species were the highest and were considered alarming. For other heavy metals (Zn, Fe and Cu) concentration were much lower for the fish species, which is also supported by other research (Losasso *et al.,* 2015). The assessment of heavy metals concentrations in fish muscle in different studies around the world is shown in Table 2. From Table 2, it is clear that the concentrations of Zn and Cu from the present study were higher than in the *Lates calcarifer* of Malaysia. The major hazardous heavy metals (Pb, Cd) concentration were lower in this study as compared to the mentioned species of different regions. The Cd content in *Fenneropenaeus indicus* of Tok Bali Port was much lower than in the species found in Gangha River, India.

Comparisons of heavy metals in various organs of fish species

The bioaccumulation of heavy metals in fish organs may differ due to the physiological differences in the function and structure of different tissues (Kotze, 1997). The gill tissue was selected for study because it is considered to be the organ most exposed to heavy metals while the liver tissue was selected because it is considered as the storage of heavy metals in the fish body and the muscle tissue was selected as it is the most popular the edible part that and hence as the main route of heavy metals exposure in humans (Agusa *et al.,* 2007; Taweel *et al.,* 2013).

Gills

The study found the highest concentration of Zn and Pb of 72.97 µg/g and 0.203 µg/g respectively were in the gills of *Leiognathus daura* species while the lowest concentrations of Zn and Pb, 0.099 µg/g and 0.014 µg/g respectively, were found in the gills

Table 2. Comparison of heavy metals in fish muscle tissue with other different studies in the world

Species	Location	Heavy metals (µg/g)					Reference
		Zn	Cu	Fe	Cd	Pb	
Euthynnus affinis	Malaysia	62.43±7.01	0.705±0.2	2.125±0.28	0.007±0.008	0.002±0.001	Present study
Pampus argenteus	Malaysia	4.827±0.67	0.038±0.01	0.038±0.02	0.004±0.001	0.024±0.04	Present study
Descapterus macrosoma	Malaysia	5.286±1.08	0.064±0.03	1.288±0.36	0.003±0.002	0.001±0.001	Present study
Leiognathus daura	Malaysia	5.456±0.67	0.018±0.004	4.309±0.68	0.004±0.003	0.003±0.002	Present study
Pampus argenteus	Hara Reserve , Iran	NA	NA	NA	0.22±0.006	0.58±0.003	Mohammadnabizadeh *et al.,* 2014
Euthynnus affinis	Karachi Fish Harbour,Pakistan	NA	NA	38.02±9.10	NA	0.46±0.14	Ahmed *et al.,* 2015
Lates calcarifer	Malaysia	6.500±1.917	0.495±0.066	5.103±1.77	0.007±0.004	0.201±0.080	Nasri *et al.,* 2017
Fenneropenaeus Indicus	Malaysia	14.4±1.58	0.34±0.39	2.11±0.87	0.04±0.03	0.008±0.01	Present study
Fenneropenaeus Indicus	Gangha, India	12-226	14-92.10	NA	NA	2.20-23.10	Bhattacharyya *et al.,* 2013

NA-Not available

of *Decapterus macrosoma*. The highest concentration were observed in the gills because, as respiratory organ, the gills extracts heavy metals along with dissolved oxygen from the water during respiratory action as reported by various authors (Zhao *et al.,* 2012, Monday *et al.,* 2007; Bashir *et al.,* 2013). The maximum concentrations of Cu, Fe and Cd were observed to be 0.79 µg/g, 3.969 µg/g and 0.191 µg/g respectively in the gills of *Euthynnus affinis* while the lowest concentration of Cu was 0.099 µg/g in the gills of *Pampus argenteus* and, the lowest concentrations of Fe, Cd and Pb were 1.313 µg/g, 0.013 µg/g and 0.014 µg/g respectively in gill of *Descapterus macrosoma*. *Pampus argenteus* is a bottom feeding species while *Euthynnus affinis* takes food from the surface waters and *Decapterus macrosoma* is pelagic (being neither close to the bottom nor near the shore) schooling species which feeds on smaller planktonic invertebrates. Differences in the trophic level of a species' habitat show different characteristics of heavy metal accumulation in fish species where the organs of fish play vital roles in binding metals but the binding abilities varies from organ to organ *(Qiao-qiao et al.,* 2007; Gu *et al.,* 2015). As *Leiognathus daura* is a demersal (it lives on or near the bottom) species, there is higher bioavailability of Zn to the species because of the availability of a higher content of Zn in the sediment and water (Akbulut and Akbulut, 2010). Bottom level inhabiting fish have higher potentials for the uptake of heavy metals than top level inhabiting fish due to direct contact with the sediment (El-Moselhy *et al.,* 2014; Wei *et al.,* 2014, Gu, 2015). The present study found the highest concentration of Zn at 72.97 µg/g in *Leiognathus daura* species while there was less Zn in *Pampus argenteus and Descapterus macrosoma.* Although, heavy metal uptake depends on various factors such as age, geographical distribution, feeding nature, trophic level and species specific factors that varied from organ to organ (El-Moselhy *et al.,* 2014).

Flesh

The determination of heavy metals in fish flesh is important as it is directly consumed by humans. The maximum level of Fe detected was detected 4.32 µg/g in the flesh of *Leiognathus daura* and the lowest amount of 0.04 µg/g was in the flesh of *Pampus argenteus*. Although both *Leiognathus daura* (blackfin ponyfish) and *Pampus argenteus* (silver pomfret) are benthopelagic species, their feeding habits are different. The maximum concentrations of Zn, Cu and Cd were observed in the flesh of *Euthynnus affinis* where the concentrations were 62.430 µg/g, 0.705 µg/g and 0.007 µg/g respectively. The minimum concentrations of Zn, Cu and Cd were 4.827 µg/g, 0.018 µg/g and 0.003 µg/g in flesh of *Pampus argenteus, Leiognathus daura* and *Descapterus macrosoma* respectively. The maximum concentration of Pb (0.024 µg/g) in flesh of *Pampus argenteus* and the minimum concentration of Pb (0.001 µg/g) was in the flesh of *Descapterus macrosoma*. The flesh of all fish species contained low amounts of Pb. The concentration of Cd was also minimum in the flesh of all fish samples studied.

Liver

Higher accumulations of Zn, Fe Cd and Pb of 43.740 µg/g, 3.897 µg/g, 1.19 µg/g and 0.41µg/g respectively were found in the liver of *Euthynnus affinis* while the lower amounts of Cd and Pb were found to be 0.114 and 0.015 µg/g in the liver of *Descapterus macrosoma*. The lowest concentration of Fe was 0.77 µg/g in the liver of *Pampus argenteus* and the lowest concentration of Zn was 9.440 µg/g in the liver of *Leiognathus daura*. However, the highest concentration of Cu was 0.637 µg/g in the liver of *Pampus argenteus* and the lowest concentration was 0.216 µg/g in the liver of *Leiognathus daura*. The liver plays major role in metabolic processes and shows a tendency to accumulate metals at higher levels (Zhao *et al.,* 2012*)*, thus a higher concentration of Cd, Pb, Zn and Fe was observed in the liver of *Euthynnus affinis*. Generally, it is considered that the liver acts as an environmental indicator of water pollution because it accumulates various kinds of pollutants at higher levels than surrounding environment and plays an important role in contaminant storage, detoxification, redistribution, transformation and act as an active pathological site (Licata *et al.,* 2005; Ebrahimpour *et al.,* 2011).

Relationships between inter metal levels in various organs and fish sizes

The relationship between metal concentrations and fish length were determined and are shown in Table 4. From the correlation (Pearson Correlation) values (r) it is clear that there is a negative correlation between fish size and metal concentration in the various organs of fish. For instance, the Fe concentration in the gills of all the species has a strong negative correlation (r=-

0.99) with the fish length and the Cd concentration in the liver of all species is also strongly negatively correlated (r= - 0.99) with fish size. The decrease in metal concentration with fish size may be due to rapid tissue growth rather than the metal uptake rate (Merciai *et al.,* 2014). From the correlation value it is seen that the metal concentration in the gills of fish has a stronger negative correlation than other organs. This is because the respiratory rates of small fish are higher than larger fish, resulting in a higher quantity of respiratory water passing through the gills and therefore higher accumulation of metals (Merciai *et al.,* 2014). Usually *Euthynnus affinis* has an average length of 45 - 60 cm, but our collected samples had an average length of 25.1 cm, and due to this small size and associated higher respiratory rate its metal accumulation in the gill was higher as compared with other species. In the case of *Leiognathus daura* the higher metal concentrations in the gills (Table 3) was observed because it is a demersal fish which lives and feeds on or near the sea floors and there is always a strong possibility of heavy metal accumulation in sediment (Bergstad, 2009). Again the high metabolic rate and faster short-term uptake by smaller species are also responsible for the higher accumulation of heavy metals (Newman and Mitz, 1988; Newman and Mitz, 1989). In our study we observed a positive correlation of Pb concentration with fish size and a negative correlation of Pb content in fish muscle with Pb content in other organs of fish.

Table 3. Heavy metal concentration (µg/g) found in different selected fish, prawn and crab species

Fish	Organ	Zn	Cu	Fe	Cd	Pb
Euthynnus affinis	gill	69.2±9.31^{aA}	0.789±0.16^{aB}	3.969±1.52^{aB}	0.191±0.15^{aB}	0.134±0.06^{aB}
	liver	43.74±8.45^{aA}	0.445±0.28^{aB}	3.897±1.55^{aB}	1.189±0.78^{aB}	0.41±0.19^{abB}
	flesh	62.43±7.01^{bA}	0.705±0.2^{aB}	2.125±0.28^{aB}	0.007±0.008^{aB}	0.002±0.001^{bB}
Pampus argenteus	gill	14.53±3.46^{aA}	0.377±0.16^{aB}	2.609±1.39^{aB}	0.03±0.01^{aB}	0.203±0.02^{aB}
	liver	15.25±4.42^{abA}	0.637±0.33^{aB}	0.77±0.12^{abB}	0.636±0.3^{bB}	0.17±0.02^{aB}
	flesh	4.827±0.67^{bA}	0.038±0.01^{bB}	0.038±0.02^{bB}	0.004±0.001^{bB}	0.024±0.04^{bB}
Descapterus	gill	13.54±1.23^{aA}	0.099±0.09^{aB}	1.313±0.25^{aB}	0.013±0.006^{aB}	0.014±0.01^{aB}
macrosoma	liver	23.69±2.75^{bA}	0.218±0.18^{aB}	1.99±0.33^{aB}	0.114±0.11^{aB}	0.015±0.009^{aB}
	flesh	5.286±1.08^{cA}	0.064±0.03^{aB}	1.288±0.36^{aB}	0.003±0.002^{aB}	0.001±0.001^{aB}
Leiognathus daura	gill	72.97±2.75^{aA}	0.338±0.13^{aB}	2.673±1.78^{aB}	0.033±0.02^{aB}	0.205±0.02^{aB}
	liver	9.44±1.08^{bA}	0.216±0.06^{abB}	1.364±1.03^{aB}	0.649±0.9^{bB}	0.15±0.08^{aB}
	flesh	5.456±0.67^{cA}	0.018±0.004^{bB}	4.309±0.68^{aB}	0.004±0.003^{bB}	0.003±0.002^{bB}
Fenneropenaeus indicus		14.4±1.58^{aA}	0.34±0.39^{aB}	2.11±0.87^{aB}	0.04±0.03^{aB}	0.008±0.01^{aB}
Portunus pelagicus		12.43±3.85^{aA}	1±0.87^{aB}	1.78±0.79^{aB}	0.09±0.07^{aB}	0.015±0.01^{aB}
WHO/FAO		150	10	43	0.2	1.5
Malaysia Food Act		100	30	NA	1	2

Letter (A, B) in rows with different letter are significantly different (P<0.05)
Letter (a, b, c) in columns with different letter are significantly different (P<0.05)
WHO: World Health Organization; FAO: Food and Agriculture Organization (FAO, 1983; MFA, 1983).

Table 4. Relationships between inter metal levels in various organs and fish sizes

		Length	Gill					Liver					Flesh				
			Zn	Cu	Fe	Cd	Pb	Zn	Cu	Fe	Cd	Pb	Zn	Cu	Fe	Cd	Pb
Length		1															
Gill	Zn	-0.75	1														
	Cu	-0.98	0.63	1													
	Fe	-0.99	0.71	0.98	1												
	Cd	-0.89	0.59	0.94	0.87	1											
	Pb	-0.50	0.42	0.39	0.55	0.07	1										
Liver	Zn	-0.57	0.22	0.69	0.54	0.86	-0.39	1									
	Cu	-0.36	-0.2	0.44	0.45	0.25	0.47	0.16	1								
	Fe	-0.63	0.49	0.69	0.59	0.89	-0.35	0.94	-0.1	1							
	Cd	-0.99	0.69	0.98	0.99	0.88	0.54	0.55	0.44	0.59	1						
	Pb	-0.98	0.63	0.99	0.98	0.95	0.38	0.70	0.44	0.70	0.98	1					
Flesh	Zn	-0.84	0.54	0.90	0.82	0.99	-0.03	0.92	0.21	0.93	0.89	0.96	1				
	Cu	-0.81	0.50	0.88	0.78	0.98	-0.08	0.94	0.21	0.94	0.72	0.89	0.99	1			
	Fe	-0.28	0.84	0.11	0.21	0.11	0.24	-0.1	-0.6	0.16	0.2	0.11	0.08	0.03	1		
	Cd	-0.95	o.64	0.98	0.94	0.99	0.24	0.79	0.34	o.80	0.94	0.99	0.96	0.95	0.13	1	
	Pb	0.06	-0.5	-0.02	0.02	-0.2	0.54	-0.3	0.85	-0.6	0.02	0.04	-0.3	-0.3	-0.64	-0.17	1

Health risk assessment

In this study both EDI and THQ were used to examine the health risk associated with the uptake of fish, prawns and crabs inhabiting in the Tok Bali coast in Kelantan, Malaysia. Health risk assessments were carried out independently for adults and children because children are considered to be more sensitive to the pollutants than adults (Shou Zhao *et al.,* 2012; Wang *et al.,* 2012).

Estimated daily intake

The EDI of heavy metals by human was evaluated according to the mean concentration of metals in aquatic organisms and the amount consumed amount. Table 5 shows the estimated daily intake of heavy metals from fish. The highest concentration of daily intake of Zn for adults was 103.86 µg/kg/day from *Euthynnus affinis* and the lowest intake was 8.03 µg/kg/day from *Pampus argenteus*. The maximum and minimum daily intake of Cu was 1.66 µg/kg/day from *Portunus pelagicus* and 0.03 µg/kg/day from *Leiognathus daura* respectively for adults while it was 1.52 and 0.03 µg/kg/day from *Portunus pelagicus* and *Leiognathus daura* respectively for children. The highest daily intake of Fe was estimated to be 7.17 µg/kg/day from *Leiognathus daura and lowest was 0.06* µg/kg/day from *Pampus argenteus* for adults. For children the highest daily intake estimated for Fe was 6.59 µg/kg/day from *Leiognathus daura* and the lowest was 0.06 µg/kg/day from *Pampus argenteus*. The daily intake of Cd of 0.15 µg/kg/day was the highest from *Portunus pelagius* and the lowest was 0.005 µg/kg/day from *Descapterus macrosoma* for adults. For children the highest daily intake was estimated to be 0.14 µg/kg/day from *Portunus pelagius* and the lowest was 0.005 µg/kg/day from *Descapterus macrosoma*. The maximum and minimum daily intake of Pb was 0.04 µg/kg/day from *Pampus argenteus* and 0.002 µg/kg/day from *Descapterus macrosoma* respectively for adults while it was 0.4 and 0.002 µg/kg/day from *Pampus argenteus* and *Descapterus macrosoma* respectively for children. All estimated daily intake values were lower than the standard estimated data (JECFA, 2009).

Target hazard quotients

All of the fish, prawn and crab species discussed in this study were easily available and commonly consumed by the people of Kelantan. The target hazard quotients is considered to recognize the non-carcinogenic health risks due to the pollutants exposure of population (Chien *et al.,* 2002; Yi *et al.,* 2017; Zhang *et al.,* 2017). In Fig. 2, there was no THQ greater than 1 through the consumption of sea food indicating that health risks associated with heavy metals exposure were insignificant. If the total target hazard quotient exceeds 1, it denotes a non-carcinogenic health risk from the accumulative effects of heavy metals among the exposed population (Yi *et al.,* 2017). The THQ of heavy metals from marine fishes, prawn and crab consumption follows the order Zn >Fe >Cd>Cu> Pb in case of adults while the THQ for children follows the order Zn >Fe > Pb>Cd>Cu. Table 5 shows the comparison of the estimated daily intake (µg/kg/day) of heavy metals from studied marine species with the recommended daily dietary allowances by Joint FAO/WHO Expert Committee on Food Additives (JECFA).

The total and individual THQ values of six metals in adults (TTHQ= 0.0012) were higher than those of

Table 5. Estimated daily intake (µg/kg/day) of heavy metals from fish

Consumer	Species	Zn	Cu	Fe	Cd	Pb
Adult	*Euthynnus affinis*	103.86	1.173	3.535	0.012	0.003
	Pampus argenteus	8.031	0.063	0.063	0.007	0.04
	Descapterus macrosoma	8.794	0.106	2.143	0.005	0.002
	Leiognathus daura	9.077	0.03	7.169	0.007	0.005
	Fenneropenaeus indicus	23.96	0.566	3.51	0.067	0.013
	Portunus pelagicus	20.68	1.664	2.96	0.15	0.025
Children	*Euthynnus affinis*	95.46	1.078	3.249	0.011	0.003
	Pampus argenteus	7.381	0.058	0.058	0.006	0.037
	Descapterus macrosoma	8.083	0.098	1.969	0.005	0.002
	Leiognathus daura	8.343	0.028	6.589	0.006	0.005
	Fenneropenaeus indicus	22.02	0.52	3.226	0.061	0.012
	Portunus pelagicus	19.01	1.529	2.722	0.138	0.023
	Tolerable limit (µg/kg/day)	300-1000	50-500	80-100	1	3.57




Fig. 2. Target hazardous quotient of heavy metals in fish and shellfish by adult and children (EA: *Euthynnus affinis*; PA: *Pampus argenteus*; DM: *Descapterus macrosoma;* LD: *Leiognathus daura;* FI: *Fenneropenaeus indicus;* PP: *Portunus pelagicus)*

children (TTHQ= 0.00103), implying that adults may suffer higher health risk (Zhao, 2012). Among all the heavy metal studied, marine fish contaminated by Zn, Fe and Cd posed relatively higher health risks, particularly for adults. Overall, the average EDI and THQ values for both adult and children were below harmful levels indicating that the consumption of both fish and shellfish around the Tok Bali Port of Kelantan, Malaysia would not result in health risks from the five studied metals.

CONCLUSION

This study revealed that the levels of heavy metals in all samples were within the acceptable limit. However, the highest levels of both essential and hazardous heavy metals were found in the gills and liver of the fish species. Among the studied metals, Cd was found to be at the highest levels followed by Pb on the basis of permissible limits. Both the EDI and THQ values revealed that the studied fish, prawns and crabs species should not be considered as a threat to consumers. The data obtained from this study can be used as a safeguard for sustaining of both native and export market of fish industries in Malaysia. In view of importance of the fish nutrition, it is necessary to maintain regular biological monitoring of the Tok Bali Port waters and fish to ensure the continuing safety of seafood. Finally safe disposal of domestic wastes and industrial effluents should be ensured and these waste should be recycled if possible to avoid their accumulation in the marine environment.

ACKNOWLEDGEMENT

This study was conducted at the Faculty of Earth Science, University Malaysia Kelantan. The authors would like to thank Rohanif and Nur Izzati, Laboratory Technologist, Environmental Monitoring Laboratory, University Malaysia Kelantan for their technical supports for analysis. The work was carried out without any financial grant from any sources.

CONFLICT OF INTEREST

The author declares that there is no conflict of interests regarding the publication of this manuscript. In addition, the ethical issues, including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, and redundancy have been completely observed by the authors.

ABBREVIATIONS

AAS	Atomic Absorption Spectrometer
WAB	Average body weight
Cd	Cadmium
^0C	Celsius
cm	Centimeter
CRM	Certified reference material
Cu	Copper
DM	*Descapterus macrosoma*
Eq.	Equation
EDI	Estimated daily intake
EA	Euthynnus affinis
ED	Exposure duration
EF	Exposure frequency
FI	Fenneropenaeus indicus
Fig.	Figure
FAO	Food and Agriculture Organization
FIR	Food ingestion rate
K	Fulton factor
g	Gram
HSD	Honestly Significant Difference
Fe	Iron
JECFA	Joint FAO/WHO Expert Committee on Food Additives
kg	Kilogram
km	Kilometer

Pb	Lead
LD	Leiognathus daura
MFA	Malaysia Food Act
m	Meter
µg/g	Microgram/gram
µg/kg	Microgram/kilogram
µg/kg/day	Microgram/kilogram/Day
PA	Pampus argenteus
r	Pearson correlation value
%	Percent
PP	Portunus pelagicus
QA	Quality assurance
QC	Quality control
RfD	Reference dose
SPSS	Statistical package for the social sciences
TA	Average exposure time for non-carcinogens
THQs	Target hazard quotients
USEPA	United States Environmental Protection Agency
WHO	World Health Organization
Zn	Zinc

REFERENCES

Agusa, T.; Kunito, T.; Sudaryanto, A.; Monirith , I.; Kan-Atireklap, S.; Iwata, H.; Ismail, A.; Sanguansin, J.; Muchtar, M.; Tana, T.S.; Tanabe, S., (2007). Exposure assessment for trace elements from consumption of marine fish in Southeast Asia. Environ. Pollut., 145(3): 766–777 **(12 pages).**

Agusa, T.; Kunito, T.; Yasunaga, G.; Iwata, H.; Subramanian, A.; Ismail, A.; Tanabe, S., (2005). Concentrations of trace elements in marine fish and its risk assessment in Malaysia. Mar. Pollut. Bull., 51(8): 896–911 **(16 pages).**

Ahmed, M.K.; Baki, M.A.; Islam, M.S.; Kundu, G.K.; Habibullah-Al-Mamun, M.; Sarkar, S.K.; Hossain, M.M., (2015). Human health risk assessment of heavy metals in tropical fish and shellfish collected from the river Buriganga, Bangladesh. Environ. Sci. Pollut. Res., 22(20): 15880–15890 **(11 pages).**

Ahmed, M.K.; Parvin, E.; Islam, M.M.; Akter, M.S.; Khan, M.S.; Al-Mamun, M.H., (2014). Lead and cadmium induced histopathological changes in gill, kidney and liver tissues of freshwater climbing perch Anabas testudineus (Bloch, 1792). Chem. Ecol., 30: 532–540 **(9 pages).**

Ahmed, Q.; Bat, L., (2015). Heavy metal levels in Euthynnus affinis (Cantor 1849) Kawakawa fish marketed at Karachi Fish Harbour, Pakistan and potential risk to human health. J. Black Sea/Mediterr. Environ., 21(1): 35-44 **(10 pages).**

Al-Busaidi, M.; Yesudhason, P.; Al-Mµghairi, S.; Al-Rahbi, W.A.; Al-Harthy, K.S.; Al-Mazrooei, N.A.; Al-Habsi, S.H., (2011). Toxic metals in commercial marine fish in Oman with reference to national and international standards. Chemosphere, 85(1): 67-73 **(7 pages).**

Amisah, S.; Obirikorang, K.A.; Boateng, D.A., (2011). Bioaccumulation of heavy metals in the Volta Clam, *Galatea paradoxa* (Born, 1778) in relation to their geoaccumulation in benthic sediments of the Volta Estuary, Ghana, Water Qual. Expo. Health., 2(3-4): 147-156 **(10 Pages).**

Anandkumar, A.; Nagarajan, R,; Prabakaran, K.; Rajaram, R., (2017). Trace metal dynamics and risk assessment in the commercially important marine shrimp species collected from the Miri coast, Sarawak, East Malaysia. Reg. Stu. Mar. Sci., 16: 79-88 **(10 pages).**

Arulkumar, A.; Paramasivam, S.; Rajaramb, R., (2017). Toxic heavy metals in commercially important food fishes collected from Palk Bay, Southeastern India. Mar. Pollut. Bull., 119(1): 454-459 **(6 pages).**

Asegbeloyin, J.N.; Onyimonyi, A.E.; Ujam, O.T.; Ukwueze, N.N.; Ukoha, P.O., (2010). Assessment of toxic trace metals in selected fish species and parts of domestic animals. Pak. J. Nutr., 9(3): 213-215 **(3 pages).**

Bashir, F.H.; Othman, M.S.; Mazlan, A.G.; Rahim, S.M.; Simon, K.D., (2013). Heavy Metal Concentration in Fishes from the Coastal Waters of Kapar and Mersing, Malaysia. Turk. J. Fish. Aqu. Sci., 13(2): 375-382 **(8 pages).**

Bergstad, O.A., (2009), Fish: Demersal Fish (Life Histories, Behavior, Adaptations), Encycl. Ocean Sci. (Second Edition), pp 458–466 **(9 pages).**

Bhattacharyya, S.B.; Roychowdhury, G.; Zaman, S.; Raha, A.K.; Chakraborty, Bhattacharjee, A.S.; Mitra, A., (2013). Bioaccumulation of Heavy Metals in Indian White Shrimp (Fenneropenaeus Indicus): A Time Series Analysis. Int. J. Life Sc. Biotechnol. Pharm. Res., 2(2): 97-113 **(17 pages).**

Chien, L.C.; Hung, T.C.; Choang, K.Y.; Yeh, C.Y.; Meng, P.J.; Shieh, M.J.; Ha, B.C., (2002). Daily intake of TBT, Cu, Zn, Cd and As for fishermen in Taiwan. Sci. Tot. Env., 285(1-3): 177–185 **(9 pages).**

Colla, N.S.L.; Botté, S.E.; Marcovecchio, J.E., (2017). Tracing Cr, Pb, Fe and Mn occurrence in the Bahía Blanca estuary throµgh commercial fish species, Chemosphere. 175: 286-293 **(8 pages).**

Ebrahimpour, M.; Pourkhabbaz, A.; Baramaki, R.; Babaei, H.; Rezaei, M., (2011). Bioaccumulation of Heavy Metals in Freshwater Fish Species, Anzali, Iran. Bull. Environ. Cont. Toxicol., 87: 386–392 **(7 pages).**

El Nemr, A .; El-Said, G.F.; Ragab, S .; Khaled, A .; El-Sikaily, A., (2016). The distribution, contamination and risk assessment of heavy metals in sediment and shellfish from the Red Sea coast, Egypt. Chemosphere, 165: 369-380 **(12 pages).**

El-Moselhy, K.M.; Othman, A.I.; Azem. H.A.E.; El-Metwally, M.E.A., (2014). Bioaccumulation of heavy metals in some tissues of fish in the Red Sea, Egypt. Egypt. J. B. App. Sci., 1(2): 97-105 **(9 pages).**

FAO, (1983). The state of food and agriculture.

Gale, N.L.; Adams, C.D.; Wixson, B.G.; Loftin, K.A.; Huang, Y.W., (2004). Lead, zinc, copper, and cadmium in fish and sediments from the Big River and Flat River Creek of Missouri's Old Lead Belt. Environ. Geochem. Health., 26(1): 37–49 **(13 pages).**

Gu, Y.G.; Lin, Q.; Huang, H.H.; Wang, L.; Ning, J.J.; Du, F.Y., (2017). Heavy metals in fish tissues/stomach contents in four marine wild commercially valuable fish species from the western continental shelf of South China Sea. Mar. Pollut. Bull., 114(2): 1125-1129 **(5 pages).**

Gu, Y.G.; Lin, Q.; Wang, X.H.; Du. F.Y.; Yu, Z.L.; Huang, H.H., (2015). Heavy metal concentrations in wild fishes captured from the South China Sea and associated health risks. Mar. Pollut. Bull., 96(1-2): 508-512 **(5 pages).**

Hajeb, P.; Jinap, S.; Ismail, A.; Fatimah, A.B.; Jamilah, B.; Rahim, M.A., (2009). Assessment of mercury level in commonly consumed marine fishes in Malaysia. Food Cont., 20(1): 79–84 **(6 pages).**

Harada, M.; Nakachi, S.; Chen, T.; Hamada, H.; Ono, Y.; Tsuda, T.; Yanagida, K.; Kizaki, T.; Ohno, H., (1999). Monitoring of mercury

pollution in Tanzinaia: relation between head hair mercury and health. Sci. Total Environ., 227(2-3): 249–256 **(8 pages).**

Hosseini, A.A.; Karbassi, A.; Hassanzadeh, B.K.; Monavari, S.M.; Sekhavatjou, M.S., (2012). Bioaccumulation of trace elements in different tissues of three commonly available fish species regarding their gender, gonadosomatic index, and condition factor in a wetland ecosystem. Environ. Monit. Assess., 184(4): 1865-1878 (**14 pages).**

Htun-Han, M., (1978). The reproductive biology of the dab Limanda limanada (L.) in the North Sea: gonadosomatic index, hepatosomatic index and condition factor. J. Fish Biol., 13(3): 351–377 **(27 pages).**

Islam, M.S.; Ahmed, M. K.; Raknuzzaman, M.; Habibullah-Al-Mamun, M.; Masunaga, S., (2014). Metal Speciation in Sediment and Their Bioaccumulation in Fish Species of Three Urban Rivers in Bangladesh. Arch. Environ. Contam. Toxicol., 68(1): 92–106 **(15 pages).**

JECFA, (2003). Joint FAO/WHO Expert Committee on Food Additives. Joint Food and Agriculture Organization/World Health Organization Expert Committee on Food Additives, Summary and Conclusions of the 61st Meeting of the Joint FAO/WHO Expert Committee on Food Additives.

JECFA, (2009). Evaluations of the Joint FAO/WHO Expert Committee on Food Additives. Joint FAO/WHO Expert Committee on Food Additives.

Jos´e, L.; Domingo, A.; Falc´o, B.G.; Llobet J. M., (2007). Benefits and risks of fish consumption Part I. A quantitative analysis of the intake of omega-3 fatty acids and chemical contaminants. Toxicol. 230(2-3): 219–226 **(8 pages).**

Kibria, K.; Hossain, M.M.; Mallick, D.; Lau, T.C.; Wu, R., (2016). Monitoring of metal pollution in waterways across Bangladesh and ecological and public health implications of pollution. Chemosphere., 165: 1-9 **(9 pages).**

Kotze, P.J., (1997). Aspects of water quality, metal contamination of sediment and fish in the Oilfants River. Disserattion, Rand Afr. University South Africa.

Lee, C.C.; Hsu, Y.C.; Kao, Y.T.; Chen, H.L., (2016). Health risk assessment of the intake of butyltin and phenyltin compounds from fish and seafood in Taiwanese population. Chemosphere, 164: 568-575 **(8 pages).**

Lee. K.; Kweon, H.Y.; Yeo, J.; Woo, S.O.; Han, S.M.; Kim, J.H., (2011). Characterization of tyrosine-rich Antheraea pernyi silk fibroin hydrolysate. Int. J. Biol. Macromol., 48(1): 223–226 **(4 pages).**

Li, J.; Sun, C.; Zheng, L.; Jiang, F.; Wang, S.; Zhuang, Z.; Wang, X., (2017). Determination of trace metals and analysis of arsenic species in tropical marine fishes from Spratly islands. Mar. Pollut. Bull., 122(1-2): 464-469 **(6 pages).**

Licata, P.; Trombetta, D.; Cristani, M.; Naccari, C.; Martino, D.; Calò, M.; Naccari, F., (2005). Heavy metals in liver and muscle of bluefin tuna (Thunnus thynnus) caught in the straits of Messina (Sicily, Italy). Environ. Monit. Assess., 107(1-3): 239248 **(10 pages).**

Linnik, P.M.; Zubenko, I.B., (2000). Role of bottom sediments in the secondary pollution of aquatic environments by heavy metal compounds: Lakes and Reservoirs. Res. Manage., 5(1): 11–21 **(11 pages).**

Losasso, C.; Bille, L.; Patuzzi, I.; Lorenzetto, M.; Binato, G.; Pozza, M.D.; Ferrè, N.; Ricci, A., (2015). Possible Influence of Natural Events on Heavy Metals Exposure from Shellfish Consumption: A Case Study in the North-East of Italy. Front Public Health., 3: 1-7 **(7 pages).**

Martín, J.R.; De Arana, C.; Ramos-Miras, J.J.; Gil, C.; Boluda, R., (2015). Impact of 70 years urban growth associated with heavy metal pollution. Environ. Pollut., 196: 156-163 **(8 pages).**

Matos, L.A.; Cunha, A.C.S.; Sousa, A.A. Maranhão, J.P.R.; Santos, N.R.S.; Gonçalves, M.M.C.; Dantas, S.M.M.M..; Sousa, J.M.C.E..; Peron, A.P.; Silva, F.C.C.D.; Alencar, M.V.O.B.; Islam, M.T.; Aguiar, R.P.S.; Melo-Cavalcante, A.A.C.; Bonecker, C.C.; Junior, H.F.J., (2017). The influence of heavy metals on toxicogenetic damage in a Brazilian tropical river. Chemosphere, 185: 852-859 **(8 pages).**

Merciai, R.; Guasch, H.; Kumar, A.; Sabater, S.; García-Berthou, E., (2014). Trace metal concentration and fish size: Variation among fish species in a Mediterranean river. Ecotox. Environ. Saf., 107: 154-161 **(8 pages).**

Merian, E., (1991). Metals and their compounds in the environment, Occurrence Analysis and Biological Relevance. Weinheim, Germany.

Mohammadnabizadeh, S.; Pourkhabbaz, A.; Afshari, R., (2014). Analysis and Determination of Trace Metals (Nickel, Cadmium, Chromium, and Lead) in Tissues of Pampus argenteus and Platycephalus indicus in the Hara Reserve, Iran. J. Toxicol., Article ID 576496, **(6 pages).**

Monday, U.E.; Nsikak, U.B., (2007). Cadmium, Copper, Lead and Zinc tissue levels in Bonga shad (Ethmalosa Fimbriata) and Tilapia (Tilapia guineensis) caught from Imo River, Nigeria. Am. J. Food Technol., 2: 45-48 **(4 pages).**

Murtala, B.A.; Abdul, W.O.; Akinyemi, A.A., (2012). Bioaccumulation of Heavy Metals in Fish (Hydrocynus Forskahlii, Hyperopisus Bebe Occidentalis and Clarias Gariepinus) Organs in Downstream Ogun Coastal Water, Nigeria. Trans. J. Agri. Sci., 4: 51-59 **(9 pages).**

MFA, (1983). Food Act. Laws of Malaysia. Act 281.

Nasri, N.N.S.; Aris. A.Z.; Mohammad, K.N.; Looi. L.J.; Kim, K.W., (2017). Bioaccumulation of Heavy Metals in Maricultured Fish, Lates calcarifer (Barramudi), Lutjanus campechanus (Red Snapper) and Lutjanus griseus (Grey Snapper). Chemosphere, 197: 318-324 **(7 pages).**

Newman, M.C.; Doubet, D.K., (1989). Size-dependence of mercury (II) accumulation kinetics in the mosquitofish, Gambusia affinis (Baird and Girard). Arch. Environ. Contam. Toxicol., 18, 819–825 **(7 pages).**

Newman, M.C.; Mitz, S.V., (1988). Size dependence of zinc elimination and uptake from water by mosquitofish Gambusia affinis (Baird and Girard). Aquat. Toxicol., 12: 17–32 **(16 pages).**

Onsanit, S.; Ke, C.; Wang, X.; Wang, K.; Wang, W., (2010). Trace elements in two marine fish cultured in fish cages in Fujian province China. Environ. Pollut., 158: 1334–1342 **(9 pages).**

Özden, Ö.; Erkan, N.; Kaplan, M. et al. , (2018). Toxic Metals and Omega3 Fatty Acids of Bluefn Tuna from Aquaculture: Health Risk and Benefits. Expo. Health., 1-10 **(10 pages).**

Part, P.; Svanberg, O.; Kiessling, A., (1985). The availability of cadmium to perfused rainbow trout gills in different water qualities. Water Res., 19: 427-434 **(8 pages).**

Perugini, M.; Visciano, P.; Manera, M. et al. , (2014). Heavy metal (As, Cd, Hg, Pb, Cu, Zn, Se) concentrations in muscle and bone of four commercial fish caught in the central Adriatic Sea, Italy. Environ. Monit. Assess., 186: 2205-2213 **(9 pages)**.

Qiao-qiao, C.; Guang-wei, Z.; Langdon, A., (2007). Bioaccumulation of heavy metals in fishes from Taihu Lake, China. J. Environ. Sci., 19(12): 1500–1504 **(5 pages).**

Salamat, N.; Movahedinia, A.; Etemadi-Deylami, E.; Mohammadi, Y., (2015). Pike (*Esox lucius*) Bio-indicator of heavy metal pollution in Anzali Wetland. Water Qual. Exposure Health., 7(2): 251-254 **(4 Pages)**.

Salleh, N.H.M.; Halim, M.A.A., (2018). Enhancing environmental sustainability over fisheries industry through proactive risk evaluation: a case of Tok Bali fishing port. J. Sustainability Sci. Manage., 4: 51-63 **(13 pages).** Sany, S.B.T.; Salleh, A.; Sulaiman,

A.H.; Sasekumar, A.; Rezayi, M.; Tehrani, G.M., (2012). Heavy metal contamination in water and sediment of the Port Klang coastal area, Selangor, Malaysia, Environ. Earth Sci., 69(6): 2013-2025 **(13 pages).**

Sarkara, T.; Alama, M.M.; Parvina, N.; Fardous, Z.; Chowdhury, A.Z.; Hossain, S.; Haque, M.E.; Biswas, N., (2016). Assessment of heavy metals contamination and human health risk in shrimp collected from different farms and rivers at Khulna-Satkhira region, Bangladesh. Toxicol., Rep. 3: 346–350 **(5 pages).**

Satarug, S.; Vesey, D.A.; Gobe, G.C., (2017). Current health risk assessment practice for dietary cadmium: Data from different countries. Food Chem. Toxicol., 106 (Part A): 430-445 **(16 pages).**

Schintu, M.; Marrucci, A.; Marras, B.; Galgani, F.; Buosi, C.; Ibba, A.; Cherchi, A., (2016). Heavy metal accumulation in surface sediments at the port of Cagliari (Sardinia, western Mediterranean): Environmental assessment using sequential extractions and benthic foraminifera, Mar. Pollut. Bull., 111(1-2): 45-56 **(12 pages).**

Sharif, A.K.M.; Mustafa, A.I.; Amin, M.N.; Safiullah, S., (1993). Trace element concentrations in tropical marine fish from the Bay of Bengal. Sci. Total Environ., 138(1-3): 223-234 **(12 pages).**

Taweel, A.; Othman, M.S.; Ahmad. A.K., (2013). Assessment of heavy metals in tilapia fish (Oreochromis niloticus) from Lagat River and Engineering Lake in Bangi, Malaysia, and evaluation of the health risk from tilapia consumption. Ecotoxicol. Environ. Saf., 93: 45-51 **(7 pages).**

USEPA, (2011). Regional screening level summary table.

Velusamy, A.; Kumar, P.S.; Ram, A.; Chinnadurai, S., (2014). Bioaccumulation of heavy metals in commercially important marine fishes from Mumbai Harbor, India. Mar. Pollut. Bull., 81(1): 218–224 **(7 pages).**

Wang, Y.; Qiao, M.; Liu, Y.; Zhu, Y., (2012). Health risk assessment of heavy metals in soils and vegetables from wastewater irrigated area, Beijing-Tianjin city cluster, China. J. Environ., Sci. 24(4): 690–698 **(9 pages).**

Wani, A.L.; Ara, A.; Usmani, J.A., (2015). Lead toxicity: a review. Interdiscip Toxicol., 8(2): 55–64 **(10 Pages).**

Wei, Y.H.; Zhang, J.Y.; Zhang, D.W.; Tu, T.H.; Luo, L.G., (2014). Metal concentrations in various fish organs of different fish species from Poyang Lake, China. Ecotoxicol. Environ. Saf., 104: 182–188 **(7 pages).**

Yi, Y.; Tang, C.; Yi, T.; Yang, Z.; Zhang, S., (2017). Health risk assessment of heavy metals in fish and accumulation patterns in food web in the upper Yangtze River, China. Ecotoxicol. Environ., Saf. 145:295–302 **(8 pages).**

Zhang, C.; Qiao, Q.; Piper, J.D.; Huang, B., (2011). Assessment of heavy metal pollution from a Fe-smelting plant in urban river sediments using environmental magnetic and geochemical methods. Environ. Pollut., 159(10): 3057-3070 **(14 pages).**

Zhang H, Cui B, Xiao R, Zhao H (2010). Heavy metals in water, soils and plants in riparian wetlands in the Pearl River Estuary, South China. Procedia. Environ. Sci., 2: 1344–1354 **(11 pages).**

Zhang, H.D.; Huang, B.; Dong, L.; Hu, W.; Akhtar, M.S.; Qu, M., (2017). Accumulation, sources and health risks of trace metals in elevated geochemical background soils used for greenhouse vegetable production in southwestern China. Ecotoxicol. Environ. Saf., 137: 233–239 **(7 pages).**

Zhao, S.; Feng, C.; Quan, W.; Chen, X.; Niu, J.; Shen, Z., (2012). Role of living environments in the accumulation characteristics of heavy metals in fishes and crabs in the Yangtze River Estuary, China. Mar. Pollut. Bull., 64(6): 1163–1171 (**9 pages).**

AUTHOR (S) BIOSKETCHES

Salam, M.A., Ph.D., Assistant Professor, Department of Environmental Science and Disaster Management, Noakhali Science and Technology University, Noakhali, Sonapur -3814, Bangladesh and Faculty of Earth Science, University, University Malaysia Kelantan, Jeli Campus, 17600 Jeli, Malaysia. Email: *s_salam1978@yahoo.com*

Paul, S.C., M.Sc., Department of Applied Chemistry and Chemical Engineering, Noakhali Science and Technology University, Noakhali, Sonapur -3814, Bangladesh. Email: *shujitchandrapaul@gmail.com*

Noor, S.N.B.M., B.Sc., Faculty of Earth Science, University, University Malaysia Kelantan, Jeli Campus, 17600 Jeli, Malaysia. Email: *sitinurainoor29@yahoo.com*

Siddiqua, S.A., M.Sc., Department of Animal Breeding and Genetics, Bangladesh Agricultural University, Mymensingh- 2202, Bangladesh. Email: *rai.boundary@gmail.com*

Aka, T.D., M.Sc., Department of Pharmacy, Noakhali Science and Technology University, Noakhali, Sonapur -3814, Bangladesh. Email: *akadasnstu@gmail.com*

Wahab, R., Ph.D., Professor, School of Engineering and Technology, University College of Technology Sarawak, Malaysia. Email: *drrazakw5181@ucts.edu.my*

Aweng,.E.R., Ph.D., Associate Professor, Faculty of Earth Science, University of Malaysia Kelantan, Jeli Campus, 17600 Jeli Malaysia. Email: *aweng@umk.edu.my*

COPYRIGHTS

Copyright for this article is retained by the author(s), with publication rights granted to the GJESM Journal. This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0/).

HOW TO CITE THIS ARTICLE

Salam, M.A.; Paul, S.C.; Noor, S.N.B.M.; Siddiqua, S.A.; Aka, T.D.; Wahab, R.; Aweng, E.R., (2019). Contamination profile of heavy metals in marine fish and shellfish. Global J. Environ. Sci. Manage., 5(2): 225-236.

DOI: 10.22034/gjesm.2019.02.08

url: https://www.gjesm.net/article_34370.html