

ORIGINAL RESEARCH PAPER

Phenol biodegradation by bacterial strain O-CH1 isolated from seashore

M. Samimi^{1,*}, M. Shahriari Moghadam²

¹Department of Chemical Engineering, Faculty of Energy, Kermanshah University of Technology, Kermanshah, Iran

²Department of Environmental Sciences, Faculty of Natural Resources, University of Zabol, Zabol, Iran

ARTICLE INFO

Article History:

Received 07 June 2019

Revised 09 September 2019

Accepted 11 November 2019

Keywords:

Biodegradation

Bioremediation

Optimization

Phenol

Taguchi

ABSTRACT

Phenol and phenolic compounds are among the most recognized environmental pollutants which exist in industrial wastewater and enter the biological cycles due to the solubility in water. Bioremediation is one of the cost-effective and Eco-friendly methods for phenol removal. In this study, the most effective phenol-degrading bacterial strain was isolated and identified from the shores of the Oman Sea by 16S rDNA. The optimal conditions of various factors, such as pH, temperature, carbon to nitrogen ratio and salinity for the phenol biodegradation, were determined using the experimental design based on Taguchi method with L₉ array (3⁴). Ability of the isolated strain (*Halomonas elongata* strain O-CH1) in degradation of different phenol concentrations was analyzed. The optimum operating conditions for phenol removal were determined in pH value of 8, temperature of 35 °C, carbon to nitrogen ratio of 100:30 (g/L) and salinity of 35 (g/L). In these conditions, 97% of the phenol was removed from the mediums. According to the optimization results, salinity and pH were the most influential factors in the biodegradation of phenol. The O-CH1 was able to grow and degrade phenol at concentrations of 250 mg/L to 1500 mg/L. Considering the high potential of this strain for phenol degradation, determining the optimal conditions for the biodegradation and its efficacy at high concentrations of phenol, the findings in this study can be used in the biological treatment of phenolic wastewater.

DOI: [10.22034/gjesm.2020.01.09](https://doi.org/10.22034/gjesm.2020.01.09)

©2020 GJESM. All rights reserved.



NUMBER OF REFERENCES

44



NUMBER OF FIGURES

4



NUMBER OF TABLES

3

*Corresponding Author:

Email: m.samimi@kut.ac.ir

Phone: +98 83 38305001

Fax: +98 83 38305006

Note: Discussion period for this manuscript open until April 1, 2020 on GJESM website at the "Show Article."

INTRODUCTION

Nowadays, environmental pollution has become one of the major concerns of societies with the increase industrial growth. Aromatic compounds are environmental pollutants that exist in different regions such as freshwater, sea and land (Deng et al., 2018). Phenol and phenolic derivatives are among the most well-known aromatic compounds that are hazardous due to teratogenic, toxic, carcinogenic and mutagenic properties, and their removal from the environment is important (Rucká et al., 2017; Youssef et al., 2019). Some of the phenol removal methods are optical oxidation, coagulation, adsorption, chemical oxidation, solvent extraction, and biodegradation. Biodegradation is an affordable method with less harmful effects on the environment compared to other methods. In this method, in addition to removing the pollutant, no secondary harmful compounds are produced (Ren et al., 2017). The ability of phenol-degrading microorganisms has been evaluated in various studies. The microorganisms which have been isolated and purified from different regions are as; *Pseudomonas putida* (Kumar et al., 2005), *Candida aquatextoris* (Jiang et al., 2015), *Pseudomonas aeruginosa* (Hasan et al., 2015), *Thauera phenolivorans* (Yin et al., 2017), *Citrobacter* sp. (Deng et al., 2018), and *Pseudomonas* sp. ATR208 (Sepehr et al., 2019). The efficiency of biodegradation depends on several factors, such as nutrients, pH, concentration of the pollutant, and temperature (Margesin et al., 2001; Mohajeri et al., 2010; Hossen et al., 2019). In order to achieve the best performance in biodegradation, the proper organism should be used in optimum conditions. For this purpose, it is necessary to optimize the physical and chemical conditions in which the organism performs the biodegradation (Sadeghi Haddad Zavareh et al., 2016). The traditional methods for determining the optimal conditions, including one-factor at any time, are time consuming and costly. These methods are not capable of examining the simultaneous effects of factors on response, and since the total number of experiments is so high, it is not economically feasible to check them all. Therefore, it is necessary to design the experiments, in which a fraction of the tests is performed using statistical methods. An experimental design based on different models and statistical methods makes it possible to examine different factors simultaneously, in addition to conducting fewer experiments (Samimi and Shahriari

Moghadam 2018). Optimization of biodegradation using Taguchi method has been evaluated in many studies such as: optimization of polycyclic aromatic compounds by *Sphingomonas* sp. (Chen et al., 2008), optimal biodegradation of 4-chlorophenol by *Candida tropicalis* PHB5 (Basak et al., 2013), optimization of crude oil biodegradation by *Marinobacter* sp. (Shahriari Moghadam et al., 2014), and optimization of biological treatment of chlorpyrifos contaminated coils (Pant and Rai, 2018). Isolation of efficient bacteria, as well as determination of optimal conditions for their performance lead to achieving high efficiency in biodegradation. The main objective of the present study was to isolate and identify most effective phenol-degrading bacterial strain from the shores of the Oman Sea in Chabahar city, Iran. The optimal conditions for various factors involved in phenol biodegradation by isolated bacterial strain were determined using the experimental design based on Taguchi method. Moreover, the ability of the purified strain in degradation of different phenol concentrations was analyzed. This study has been carried out in the Research Laboratory of Kermanshah University of Technology and Zabol University, Iran in 2019.

MATERIALS AND METHODS

Isolation of phenol-degrading bacteria and its ability in phenol biodegradation

Aerobic layer sediments along with seawater from different parts of the coast of Chabahar city were randomly sampled in sterile bottles, stored at 4 °C and transferred to the laboratory. 5 g of the sediment samples was added to 50 ml of the mineral salt medium (MSM) containing phenol as a carbon source (200 mg/L). The MSM contains 0.5g K_2HPO_4 , 1g NH_4Cl , 0.01g $FeSO_4 \cdot 7H_2O$ as well as 1 ml of micro elements in 1000 ml of sea water. The solution of micro elements contains 70mg $ZnCl_2$, 100 mg $MnCl_2 \cdot 4H_2O$, 200mg $CoCl_2 \cdot 6H_2O$, 100mg $NiCl_2 \cdot 6H_2O$, 20mg $CuCl_2 \cdot 2H_2O$, 50mg $NaMoO_4 \cdot 2H_2O$, 26mg $Na_2SeO_3 \cdot 5H_2O$, 10mg $NaVO_3 \cdot H_2O$, 30mg $Na_2WO_4 \cdot 2H_2O$, 1ml HCl (25%) in 1000 ml distilled water (Schlegel, 1992). After observing the growth in the medium, 5 ml of the medium was added to the new conical flask containing 50 ml of fresh MSM, and the process was repeated 3 times. After serial dilution, 1 ml of the medium was cultured on plates containing nutrient agar medium. Finally, different strains were

purified using the serial dilution method. The ability of isolated bacterial strains to biodegrade phenol was also evaluated. For this purpose, the bacteria were inoculated first into the nutrient broth medium and incubated for 24 hours. After centrifugation, the bacteria were used for inoculation in the flasks containing MSM along with phenol.

Identification of isolated strain

Identification of isolated strain was carried out using 16S rDNA sequencing. First the purified strain was cultured in a nutrient agar medium. Genomic DNA was extracted by DNA extraction kit (Roche, Germany). In order to determine the concentration of DNA and its purity, DNA absorption was measured at 260 and 280 nm. The proliferation of the 16S rDNA was performed using the primers 27F: 5-AGA GTT TGA TCC TGG CTC AG-3 and 1510R: 5-GGT TAC CTT GTT ACG ACT T-3 (Sadeghi Haddad Zavareh *et al.*, 2016) by Primus 25 advanced® thermocycler, and the nucleotide sequence was determined by Takapu-Zist Gene Molecular Biotechnology Co., Iran. After determining the desired sequences, homology searches were performed using the BLAST database (Shahriari Moghadam *et al.*, 2016).

Determination of the optimal conditions for phenol biodegradation

The optimal conditions for the phenol biodegra-

tion by isolated bacterial strain were determined by the experimental design based on Taguchi method with L_9 array (3⁴). The optimal conditions for the main operational factors involved in phenol biodegradation such as pH, temperature, carbon to nitrogen ratio and salinity were evaluated. Table 1 shows the range of levels of effective factors, and design of the experiments for the variables based on L_9 array (3⁴) is presented in Table 2. Experiments were performed in 250 ml sterile flasks in a shaker at 140 rpm. To reach a number of bacteria required for each treatment, the purified strain was proliferated in a nutrient broth medium. After centrifuging the medium, the saline serum was added, and to reach the final OD of 0.1, the bacteria were inoculated to 50 ml of MSM containing 200 mg/L of phenol. After 4 days, the phenol removal by bacterial strain was measured in different treatments. Non-inoculated culture medium was used for non-biological removal. All the experiments were performed in 3 replicates.

Effect of phenol concentration on bacterial growth and phenol degradation

The O-CH1 was inoculated in a conical flask containing 50 ml of MSM and exposed to different amounts of phenol (250, 500, 750, 1000, 1250, 1500 mg/L) as the only source of carbon and energy. The amounts of phenol removal and bacterial growth (optical density) were measured every 12 hours.

Table 1: The range of levels of various factors

Factors	Symbol	Range of levels		
		1	2	3
pH	A	6	7	8
Temperature (°C)	B	20	25	35
Carbon to nitrogen ratio (g/L)	C	100:10	100:20	100:30
Salinity (g/L)	D	15	25	35

Table 2: Design of experiments for independent variables based on L_9 array (3⁴)

Run No.	Manipulated variables			
	A	B	C	D
1	1	1	1	1
2	2	2	2	1
3	3	3	3	1
4	1	3	2	2
5	2	1	3	2
6	3	2	1	2
7	1	2	3	3
8	2	3	1	3
9	3	1	2	3

All experiments were carried out according to the optimal conditions obtained in the previous steps, and the flasks were inoculated in an orbital shaker at 140 rpm.

Measurement of phenol degradation

In this study, phenol concentration was measured by the colorimetric method. Phenolic samples were coupled with 4-amino anti-pyrene, and oxidation in alkaline conditions was carried out using potassium ferricyanide. The red color produced in this reaction was measured at 500 nm. The standard curve was used to measure the amount of phenol remaining in the solution (APHA, 1998).

Statistical Methods

The Statistica 8.0 software was used to design the experiments based on the Taguchi method (Montgomery, 2017). The factorial ANOVA of the design module was used to identify the factors affecting the amount of phenol biodegradation.

RESULTS AND DISCUSSION

Four bacterial strains, capable of phenol biodegradation, were isolated from the sediments and water of the coastal area of Chabahar, Oman Sea. The results showed that the bacterial consortium

had a better ability to biodegrade phenol than purified strains, so that almost the whole phenol in the inoculated medium was degraded by the bacterial consortium at the end of the experiment. Various studies have shown that biodegradation by bacterial consortium is more efficient than pure bacterial strain. This could be due to the greater ability of the enzyme as well as the co-metabolism processes of bacterial consortium to produce the pure bacterial strains (Sugiura *et al.*, 1996; Casellas *et al.*, 1998). Identification of the most effective purified strain with the ability to biodegrade phenol, called O-CH1, showed that the strain was 98% similar to *Halomonas elongata*. *Halomonas* sp. includes some species that have a great ability to decompose various hydrocarbons. Haddadi and Shavandi (2013) reported that *Halomonas* sp. strain PH2-2 had the ability to biodegrade phenol to a concentration of 1100 mg/L. Mnif *et al.*, (2009) identified a new strain of *Halomonas* sp. capable of degradation hexadecane, crude oil, motor oil and diesel fuel at a salt concentration of 10%. Other studies have also shown that *H. campisalis* uses benzoate, salicylic acid, *H. eurihalina* from phenol and crude oil as the only source of carbon and energy (Peytone *et al.*, 2007; Calvo *et al.*, 2002). In addition, phenol biodegradation by *H. organivorans* (Bonfá *et al.*, 2013) and *Halomonas anticariensis*




Fig. 1: Phenol degradation percentage by strain O-CH1 in different treatments

FP35 (Tena-Garitaonaindia *et al.*, 2019) has been reported. The results of phenol biodegradation in different treatments showed that the highest and lowest phenol degradation percentages were obtained in treatments No. 8 and No. 1, respectively (Fig. 1).

According to the ANOVA results shown in Table 3, all the studied factors had a significant effect ($P < 0.05$) on phenol biodegradation by strain O-CH1.

The results of different levels of factors affecting the phenol biodegradation by strain O-CH1 are shown in Fig. 2. Among the factors studied based on effect size, salinity had the greatest effect (1.47) and pH (1.12), temperature (0.96) and carbon to nitrogen ratio (0.76) ranked the second, third and fourth parameters which were effective in phenol biodegradation respectively.

In different studies, Taguchi method has been used to optimize biodegradation of various compounds such as: optimization of biodegradation of phenol-contaminated wastewater (Hsien *et*

al., 2005), optimization of crude biodegradation by *Marinobacter* sp. (Shahriari Moghadam *et al.*, 2014), optimization of the factors affecting Malachite Green's biodegradation (Daneshvar *et al.*, 2007), optimization of phenol degradation by *Alcaligenes faecalis* (Kumar *et al.*, 2013) and optimization of phenol biodegradation by *Candida tropicalis* (Basak *et al.*, 2013). In the present study, Taguchi method was used to determine the optimal conditions for the phenol biodegradation by strain O-CH1. The analyses revealed that the optimum operating conditions for the phenol removal were pH value of 8, temperature of 35 °C, carbon to nitrogen ratio of 100:30 (g/L) and salinity of 35 (g/L). Under these conditions, 97% of phenol in the medium was degraded by strain O-CH1. The salinity, which has the greatest impact on the amount of phenol biodegradation, depends on environmental conditions such as sediment type, bacterial population, type of compound, etc. Salinity not only affects the metabolism of degrading bacteria,

Table 3: ANOVA analysis for the phenol biodegradation by strain O-CH1

Factors	p – value	F – value	Sum of squares
A	0.00	27.84	55.91
B	0.00	15.02	30.16
C	0.03	3.96	7.96
D	0.00	35.91	71.82




Fig. 2: Effect of different levels of factors on the phenol biodegradation by strain O-CH1

but also affects the solubility of cyclic compounds (Tremblay et al., 2005). Another factor affecting biodegradation is pH of the environment, which was determined to be 8 for optimum conditions for phenol degradation by O-CH1 strain in this study. The pH of the environment leads to changes in bacterial growth, enzymatic activities, intracellular transport, and the degree of solubility of food (Lin et al., 2010). In marine environments, the pH is often constant and partly alkaline. Generally, the optimum pH varies according to the type of microorganism. Some bacteria, such as Mycobacterium sp., perform better at acidic pH due to the greater penetration of cyclic compounds into them, whereas Pseudomonas

genus prefers neutral pH for organic matter decomposition (Kim and Freeman, 2005). Other studies have shown that *Rhodococcus pyridinivorans* shows the best biodegradation performance of phenol at pH 8 (Shahriari Moghadam et al., 2016). The results of determining optimum conditions of phenol biodegradation showed that the highest amount of phenol removal was obtained at 35 °C. In general, temperature plays an important role in controlling the properties and rate of hydrocarbon metabolism. Temperature changes the population and structure of bacteria, the rate of metabolism, and the physical and chemical properties of pollutants (Atlas, 1981). Previous studies have




Fig. 3: Bacterial growth of strain O-CH1 in different concentrations of phenol




Fig. 4: Percentage of phenol degradation by strain O-CH1 in different concentrations of phenol

shown that most of the mesophilic bacteria have the highest efficiency in decomposing petroleum hydrocarbons in the temperature range of 30 to 40 °C (Leahy and Colwell 1990). Similar results were obtained for the biodegradation of phenanthrene in river sediments (Yuan *et al.*, 2001). The results obtained in the present study confirm the results reported in the above-mentioned studies. Nutrients are one of the most important factors in the biodegradation of cyclic compounds, and nutrient deficiencies decrease the number and efficiency of bacteria. In bioremediation, it is important to determine the optimum nutrient concentration to reduce the environmental damage (Bao *et al.*, 2012). In the present study, 100:30 carbon to nitrogen ratio was determined for the efficient decomposition of phenol. Different bacteria at different concentrations of nitrogen source have their own optimal performances and it is important to determine the lowest concentration of nitrogen source where the bacteria perform best. Leys *et al.* (2005) showed that a low ratio of nitrogen to carbon decreases the bacterial growth. The results obtained from determination of optimal conditions of phenol consumption by strain O-CH1 also confirm the above results. In fact, the rate of biodegradation increased with the increase of the nitrogen source. In general, the rate of substrate degradation depends on the substrate concentration and the type of substrate degrading organisms (Okpokwasili and Nweke, 2006). Strain O-CH1 was able to grow and degrade at various concentrations of phenol (Figs. 3 and 4). Growth rate was increased with the increase of phenol concentration up to 750 mg/L, however at higher concentrations, as the lag phase and the substrate inhibitory effect increased, the growth rate was decreased. Other studies have also shown the increase of phenol percentage with the increase of substrate concentration (Shahriari Moghadam *et al.*, 2016), which confirms the results obtained in the present work.

CONCLUSION

In the present study, *Halomonas elongata* O-CH1 as the most effective phenol-degrading bacterial strain, was isolated and purified from the Oman Sea-Chabahar coast. Compared to purified strains, bacterial consortium had a better ability to biodegrade phenol, so that almost the whole

phenol in the inoculated medium was degraded by the bacterial consortium. All the experimental factors had a significant effect on the rate of phenol biodegradation by strain O-CH1. Among the factors selected based on the rate of effect, salinity and pH were the most important factors in the biodegradation of phenol. Optimum operating conditions for phenol removal were determined as pH value of 8, temperature of 35 °C, carbon to nitrogen ratio of 100:30 (g/L) and salinity of 35 (g/L) based on Taguchi method with L_9 array (3^4). Under optimal conditions, more than 97% of the phenol was degraded. Given the high potency of purified strain in phenol degradation and determination of optimum conditions as well as its contribution to bacterial growth and biodegradation in high amounts of phenol, the results of this study can be used in the bioremediation of the water contaminated with phenolic compounds.

ACKNOWLEDGMENTS

The current study was financially supported by Kermanshah University of Technology [grant No. S/P/T 1192]. The authors are also thankful to the authorities of Zabol University, whom provided the laboratory equipment.

CONFLICT OF INTEREST

The author declares that there is no conflict of interests regarding the publication of this manuscript. In addition, the ethical issues, including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, and redundancy have been completely observed by the authors.

ABBREVIATIONS

A	pH
ANOVA	Analysis of variance
B	Temperature
C	Carbon to nitrogen ratio
$CoCl_2 \cdot 6H_2O$	Cobaltous chloride hexahydrate
$CuCl_2 \cdot 2H_2O$	Cupric chloride dihydrate
D	Salinity
$FeSO_4$	Ferrous sulfate

<i>Fig.</i>	Figure
<i>g</i>	Gram
<i>h</i>	Hour
K_2HPO_4	Dipotassium phosphate
<i>l</i>	Liter
<i>mg</i>	Milligram
<i>ml</i>	Millilitre
$MnCl_2 \cdot 4H_2O$	Manganese (II) chloride tetrahydrate
$NaMoO_4 \cdot 2H_2O$	Molybdc acid sodium salt dihydrate
$Na_2SeO_3 \cdot 5H_2O$	Disodium selenite (IV) pentahydrate
$NaVO_3 \cdot H_2O$	Sodium metavanadate hydrate
$Na_2WO_4 \cdot 2H_2O$	Sodium tungstate dihydrate
NH_4Cl	Ammonium chloride
$NiCl_2 \cdot 6H_2O$	Nickel(II) chloride hexahydrate
<i>nm</i>	Nanometer
<i>No</i>	Number
<i>OD</i>	Optical Density
<i>rpm</i>	Revolutions per minut
$ZnCl_2$	Zinc chloride
$^{\circ}C$	Celsius
%	Percentage

REFERENCES

- APHA., (1998). Standard methods for the examination of water and wastewater. Joint publication of the American Public Health Association, the American Water Works Association and the Water Environment Federation.
- Atlas, R.M., (1981). Microbial degradation of petroleum hydrocarbons. An environmental perspective. *Microbiol Rev.*, 45(1): 180-209 (30 pages).
- Bao, M.T.; Wang, L.N.; Sun, P.Y.; Cao, L.X.; Zou, J.; Li, Y.M., (2012). Biodegradation of crude oil using an efficient microbial consortium in a simulated marine environment. *Mar. Pollut. Bull.*, 64(6): 1177-1185 (9 pages).
- Basak, B., Bhunia, B., Dutta, S. and Dey, A., (2013). Enhanced biodegradation of 4-chlorophenol by *Candida tropicalis* PHB5 via optimization of physicochemical parameters using Taguchi orthogonal array approach. *Int. Biodeterior Biodegrad.*, 78: 17-23 (7 pages).
- Basak, B.; Bhunia, B.; Mukherjee, S.; Dey, A., (2013). Optimization of physicochemical parameters for phenol biodegradation by *Candida tropicalis* PHB5 using Taguchi methodology. *Desalin. Water Treat. J.*, 51(34-36): 6846-6862 (17 pages).
- Bonfá, M.R.L.; Grossman, M.J.; Piubeli, F.; Mellado, E.; Durrant, L.R., (2013). Phenol degradation by halophilic bacteria isolated from hypersaline environments. *Biodegradation.*, 24(5): 699-709 (11 pages).
- Calvo, C.; Martinez-checa, F.; Toledo, F.L.; Porcel, J.; Quesada, E., (2002). Characteristics of bioemulsifiers synthesised in crude oil media by *Halomonas eurihalina* and their effectiveness in the isolation of bacteria able to grow in the presence of hydrocarbons. *Appl. Microbiol. Biotechnol.*, 60(3): 347-351 (5 pages).
- Casellas, M.; Grifoll, M.; Sebaste, J.; Solanas, A. M.; (1998). Isolation and characterization of a 9-fluorenone-degrading bacterial strain and its role in synergistic degradation of fluorene by a consortium. *Can. J. Microbiol.*, 44(8): 734-742 (9 pages).
- Chen, J.; Wong, M.H.; Wong, Y.S.; Tam, N.F.Y., (2008). Multi-factors on biodegradation kinetics of polycyclic aromatic hydrocarbons (PAHs) by *Sphingomonas* sp., a bacterial strain isolated from mangrove sediment. *Mar. Pollut. Bull.*, 57(6-12): 695-702 (8 pages).
- Daneshvar, N.; Khataee, A.R.; Rasoulifard, M.H.; Pourhassan, M., (2007). Biodegradation of dye solution containing Malachite Green: Optimization of effective parameters using Taguchi method. *J. Hazard. Mater.*, 143(1-2): 214-219 (6 pages).
- Deng, T.; Wang, H.; Yang, K., (2018). Phenol biodegradation by isolated *Citrobacter* strain under hypersaline conditions. *Water Sci. Technol.*, 77(2): 504-510 (7 pages).
- Haddadi, A.; Shavandi, M., (2013). Biodegradation of phenol in hypersaline conditions by *Halomonas* sp. strain PH2-2 isolated from saline soil. *Int. Biodeterior Biodegrad.*, 85: 29-34 (6 pages).
- Hasan, S.A.; Jabeen, S., (2015). Degradation kinetics and pathway of phenol by *Pseudomonas* and *Bacillus* species. *Biotechnol Biotechnol Equip.*, 29(1): 45-53 (9 pages).
- Hossen, M.Z.; Hussain, M.E.; Hakim, A.; Islam, K.; Uddin, M.N.; Azad, A.K., 2019. Biodegradation of reactive textile dye Novacron Super Black G by free cells of newly isolated *Alcaligenes faecalis* AZ26 and *Bacillus* spp obtained from textile effluents. *Heliyon.*, 5(7), e02068 (30 pages).
- Hsien T.Y.; Lin, Y.H., (2005). Biodegradation of phenolic wastewater in a fixed biofilm reactor, *Biochem. Eng. J.*, 27(2): 95-103 (9 pages).
- Jiang, Y.; Yang, K.; Wang, H.; Shang, Y.; Yang, X., (2015). Characteristics of phenol degradation in saline conditions of a halophilic strain JS3 isolated from industrial activated sludge. *Mar. Pollut. Bull.*, 99(1-2): 230-234 (5 pages).
- Kim, Y.H.; Freeman, J.P., (2005). Effects of pH on the degradation of phenanthrene and pyrene by *Mycobacterium vanbaalenii* PYR-1. *Appl. Microbiol. Biotechnol.*, 67(2): 275-285 (11 pages).
- Kumar, A.; Bhunia, B.; Dasgupta, D.; Mandal, T.; Dey, A.; Datta, S.; Bhattacharya, P., (2013). Optimization of culture

- condition for growth and phenol degradation by *Alcaligenes faecalis* JF339228 using Taguchi Methodology. *Desalin Water Treat.*, 51(16-18): 3153-3163 **(11 pages)**.
- Kumar, A.; Kumar, S.; Kumar, S., (2005). Biodegradation kinetics of phenol and catechol using *Pseudomonas putida* MTCC 1194. *Biochem. Eng. J.*, 22(2): 151-159 **(9 pages)**.
- Leahy, J.G.; Colwell, R.R., (1990). Microbial degradation of hydrocarbons in the environment. *Microbiol. Rev.*, 54(3): 305-315 **(11 pages)**.
- Leys, N.M.; Bastiaens, L.; Verstraete, W.; Springael, D., (2005). Influence of the carbon/nitrogen/phosphorus ratio on polycyclic aromatic hydrocarbon degradation by *Mycobacterium* and *Shingomonas* in soil. *Appl. Microbiol. Biotechnol.*, 66(6): 726-736 **(11 pages)**.
- Lin, C.; Gan, L.; Chen, Z. L., (2010). Biodegradation of naphthalene by strain *Bacillus fusiformis* (BFN). *J. Hazard. Mater.*, 182(1-3): 771-777 **(7 pages)**.
- Margesin, R.; Schinner, F., (2001). Biodegradation and bioremediation of hydrocarbons in extreme environments. *Appl. Microbiol. Biotechnol.*, 56(5-6): 650-663 **(14 pages)**.
- Mnif, S.; Chamkha, M.; Sayadi, S., (2009). Isolation and characterization of *Halomonas* sp. Strain C2SS100, a hydrocarbon-degrading bacterium under hypersaline conditions. *J. Appl. Microbiol.*, 107(3): 785-794 **(10 pages)**.
- Mohajeri, L.; Aziz, H.A.; Isa, M.H.; Zahed, M.A., (2010). A statistical experiment design approach for optimizing biodegradation of weathered crude oil in coastal sediments. *Bioresour. Technol.*, 101 (3): 893-900 **(8 pages)**.
- Montgomery, D.C., (2017). Design and analysis of experiments, John Wiley and sons, **(630 pages)**.
- Okpokwasili, G.C.; Nweke, C.O., (2006). Microbial growth and substrate utilization kinetics. *Afr. J. Biotechnol.*, 5(4): 305-317 **(13 pages)**.
- Pant, A.; Rai, J.P.N., (2018). Bioremediation of chlorpyrifos contaminated soil by two phase bioslurry reactor: Processes evaluation and optimization by Taguchi's design of experimental (DOE) methodology. *Ecotoxicol. Environ. Saf.*, 150: 305-311 **(7 pages)**.
- Peytone, B.M.; Oie, C.S.; Albaugh, C.E., (2007). Benzoate and salicylate degradation by *Halomonas campisalis*, an alkaliphilic and moderately halophilic microorganism. *Water Res.*, 41(6): 1235-1242 **(8 pages)**.
- Ren, L.F.; Chen, R.; Zhang, X.; Shao, J.; He, Y., (2017). Phenol biodegradation and microbial community dynamics in extractive membrane bioreactor (EMBR) for phenol-laden saline wastewater. *Bioresour. Technol.*, 244(1): 1121-1128 **(8 pages)**.
- Rucká, L.; Nešvera, J.; Pátek, M., (2017). Biodegradation of phenol and its derivatives by engineered bacteria: current knowledge and perspectives. *World J. Microbiol. Biotechnol.*, 33(9): 174 **(8 pages)**.
- Sadeghi Haddad Zavareh, M.; Ebrahimpour, G.; Sahahriari Moghadam, M.; Fakhari, J.; Abdoli, T., (2016). Bioremediation of Crude Oil Using Bacterium from the Coastal Sediments of Kish Island, Iran. *Iran. J. Public Health.* 45(5): 670-679 **(10 pages)**.
- Samimi, M.; Shahriari Moghadam, M., (2018). Optimal conditions for biological removal of ammonia from wastewater of a petrochemical plant using the response surface methodology. *Global J. Environ. Sci. Manage.*, 4(3): 315-324 **(10 pages)**.
- Sugiura, K.; Ishihara, M.; Shimauchi, T.; Harayama, S., (1996). Physicochemical properties and biodegradability of crude oil. *Environ. Sci. Technol.*, 31(1): 45-51 **(7 pages)**.
- Schlegel, H.G., (1992). *Allgemeine Mikrobiologie: Auflage, Georg Thieme Verlag (655 pages)*.
- Shahriari Moghadam, M.; Ebrahimpour, G.; Abtahi, B.; Ghassempour, A.; Hashtroudi, M.S., (2014). Biodegradation of polycyclic aromatic hydrocarbons by a bacterial consortium enriched from mangrove sediments. *J. Environ. Health Sci. Eng.*, 12(1): 1-9 **(9 pages)**.
- Shahriari Moghadam, M.; Ebrahimpour, G.; Abtahi, B.; Khazaei, N.; Karbasi, N., (2014). Statistical optimization of crude oil biodegradation by *Marinobacter* sp. isolated from Qeshm Island, Iran. *Iran J. Biotechnol.*, 12(1):35-41 **(7 pages)**.
- Shahriari Moghadam, M.; Safaei, N.; Ebrahimpour, G.H., (2016). Optimization of phenol biodegradation by efficient bacteria isolated from petrochemical effluents. *Global J. Environ. Sci. Manage.*, 2(3): 249-256 **(8 pages)**.
- Sepehr, S.; Shahnavaz, B.; Asoodeh, A.; Karrabi, M., 2019. Biodegradation of phenol by cold-tolerant bacteria isolated from alpine soils of Binaloud Mountains in Iran. *J. Environ. Sci. Health. Part A*, 54(4): 367-379 **(13 pages)**.
- Tena-Garitaonaindia, M.; Llamas, I.; Toral, L.; Sampedro, I., 2019. Chemotaxis of halophilic bacterium *Halomonas anticariensis* FP35 towards the environmental pollutants phenol and naphthalene. *Sci Total Environ.*, 669: 631-636 **(6 pages)**.
- Tremblay, L.; Kohl, S.D.; Rice, J.A.; Gagne, J.P., (2005). Effects of temperature, salinity, and dissolved humic substances on the sorption of polycyclic aromatic hydrocarbons to estuarine particles. *Mar. Chem.*, 96(1-2): 21-34 **(14 pages)**.
- Yin, Y.; Wang, Y.; Tang, W.; Song, L., (2017). *Thaueria phenolivorans* sp. nov., a phenol degrading bacterium isolated from activated sludge. *Antonie van Leeuwenhoek.*, 110(12):1681-1690 **(10 pages)**.
- Yuan, S.Y.; Chang, J.S.; Yue, J.H.; Chang, B.V., (2001). Biodegradation of phenanthrene in river sediment. *Chemosphere.*, 43(3): 273-278 **(6 pages)**.
- Youssef, M.; El-Shatoury, E.H.; Ali, S.S.; El-Taweel, G.E., 2019. Enhancement of phenol degradation by free and immobilized mixed culture of *Providencia stuartii* PL4 and *Pseudomonas aeruginosa* PDM isolated from activated sludge. *Bioremediat J.*, 1-19 **(19 pages)**.

AUTHOR (S) BIOSKETCHES

Samimi, M., Ph.D., Assistant Professor, Department of Chemical Engineering, Faculty of Energy, Kermanshah University of Technology, Kermanshah, Iran. Email: m.samimi@kut.ac.ir

Shahriari Moghadam, M., Ph.D., Associate Professor, Department of Environmental Sciences, Faculty of Natural Resources, University of Zabol, Zabol, Iran. Email: mohsen.shahriari@uoz.ac.ir

COPYRIGHTS

© 2020 The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, as long as the original authors and source are cited. No permission is required from the authors or the publishers.



HOW TO CITE THIS ARTICLE

Samimi, M.; Shahriari Moghadam, M., (2020). Phenol biodegradation by bacterial strain O-CH1 isolated from seashore. *Global J. Environ. Sci. Manage.*, 6(1): 109-118.

DOI: [10.22034/gjesm.2020.01.09](https://doi.org/10.22034/gjesm.2020.01.09)

url: https://www.gjesm.net/article_37147.html

